

What Can Inflammatory Markers Tell Us About Tachycardia in Children?

Irving Gabriel Araújo Bispo,¹ Eloisa Faggiani Santos,¹ Ana Beatriz Aparecida Rocha,¹ Millene Gonzalez Manente,¹ Marina Guerriero Morata¹

Universidade Municipal de São Caetano do Sul (USCS),¹ São Caetano do Sul, SP – Brazil

Editorial referring to the article: *The Relationship Between Inflammatory Markers and Supraventricular Tachycardia in Children*

The supraventricular tachycardia (SVT) can be described as an irregular heart rhythm with an increased heart rate originating above the bundle of His. The heart rate is > 100 bpm. This condition can be particularly distressing for children and their families.¹

The incidence estimate of SVT in children is inaccurate because symptoms are usually nonspecific, such as sweating, pallor, apathy, inappetence, and tachycardia. Furthermore, spontaneous resolution can occur in some cases. Nonetheless, studies estimate that in children without congenital heart pathologies, the incidence is 24.4 per 100,000 live births. Still, the incidence is lower than in adults, and SVT emergencies are rare.¹

Although the mechanisms behind this condition are complex, growing evidence suggests a significant role of systemic inflammation. Inflammatory markers are released in response to tissue damage, injury, or destruction caused by physical, chemical, or pathogenic factors. These markers are associated with SVT, as inflammation can alter the electrophysiological properties of the heart by disrupting the electrical channels that control the heartbeat, affecting depolarization, and interfering with cardiac remodeling. This disruption can lead to arrhythmias, including SVT.¹

Thus, given the clinical relevance of SVT and its association with inflammatory markers, the study “The Relationship Between Inflammatory Markers and Supraventricular Tachycardia in Children” aims to evaluate the correlation between inflammatory markers — mainly the monocyte/lymphocyte (MLR) and neutrophil/lymphocyte (NLR) ratios — and the occurrence of SVT.^{1,2}

To perform the analysis, the study was conducted retrospectively, comparing a control group with patients diagnosed with SVT. Both groups consisted of 33 individuals from the same age range. The blood count, electrocardiogram, echocardiogram, as well as the MLR and NLR ratios of all participants, were evaluated.²

The study compared patients with SVT to a control group and observed that patients with the arrhythmia had higher white blood cell and monocyte counts. Besides, the MLR

was significantly higher in these patients, suggesting that inflammation may be involved in SVT. No differences were observed between the groups regarding age, lymphocyte counts, hemoglobin, hematocrit, platelets, or the NLR.²

Upon analyzing the echocardiogram exams, it was observed that 10 patients with SVT had normal findings. However, some showed different results, such as mitral insufficiency (four cases), patent foramen ovale (two cases), and other minor heart problems.²

These findings highlight a possible link to inflammation and suggests that the MLR may serve as a valuable tool in assessing and potentially predicting the risk of SVT in children.

Understanding the role of inflammation in pediatric SVT is critical, as early identification and intervention can significantly improve long-term outcomes.

Early detection is one of the most important findings of the study, and the MLR, a simple blood test, could potentially assist in the early identification of children at increased risk for SVT. Additionally, improved risk stratification using this information could aid in better assessing the severity of the condition and adapting treatment plans accordingly.^{3,4}

Another important finding influencing new therapeutic approaches is that underlying inflammatory processes may offer new avenues for treating SVT in children, in addition to traditional medications.^{5,6}

However, further research is crucial. Larger studies are needed to confirm these findings and explore the impact of other inflammatory markers. Besides, investigating the underlying causes of this inflammation in children with SVT is essential. There are uncertainties, such as the existence of specific infections, autoimmune conditions, or environmental factors that contribute to this elevated inflammatory state.^{7,8}

In conclusion, the study provides compelling evidence for the involvement of inflammation in pediatric SVT. By focusing on accessible biomarkers, we can improve the diagnosis, risk stratification, and management of this important cardiac condition in children. This research highlights the critical need for continued investigation into the interaction between inflammation and heart health, particularly in children.

Keywords

Tachycardia; Inflammation; Echocardiography

Mailing Address: Irving Gabriel Araújo Bispo •

Universidade Municipal de São Caetano do Sul. Rua Santo Antônio, 50.

Postal code: 09550-051. Centro, São Caetano do Sul, SP - Brazil

E-mail: irvingbispo@yahoo.com.br

DOI: <https://doi.org/10.36660/ijcs.20250024>




Figure 1 – Comparison between control group and group with supraventricular tachycardia. SVT: Supraventricular Tachycardia; NLR: neutrophil/lymphocyte ratios.

References

1. Magalhães LP, Guimarães I, Melo SL, Mateo E, Andalaft RB, Xavier L, et al. Diretriz de Arritmias Cardíacas em Crianças e Cardiopatias Congênitas SOBRAC e DCC - CP. Arq Bras Cardiol. 2016;107(1 Suppl 3):1-58. doi: 10.5935/abc.20160103.
2. Durgut SS, Kiztanır H, Yıldız Y. The Relationship between Inflammatory Markers and Supraventricular Tachycardia in Children. Int J Cardiovasc Sci. 2025; 38:e20240151. doi: 10.36660/ijcs.20240151.
3. Endres KM, Kierys K, Shang Y, Zhou S, Ceneviva GD, Thomas NJ, et al. A Multicenter Retrospective Evaluation of Specialized Laboratory Investigations in the Workup of Pediatric Patients with New-Onset Supraventricular Tachycardia. J Emerg Nurs. 2022;48(6):678-87.e1. doi: 10.1016/j.jen.2022.07.002.
4. Helton MR. Diagnosis and Management of Common Types of Supraventricular Tachycardia. Am Fam Physician. 2015;92(9):793-800.
5. Ashok A, Cabalag M, Taylor DM. Usefulness of Laboratory and Radiological Investigations in the Management of Supraventricular Tachycardia. Emerg Med Australas. 2017;29(4):394-9. doi: 10.1111/1742-6723.12766.
6. Szydłowski L, Skierska A, Markiewicz-Loskot G, Mazurek B, Morka A, Undas A. The Role of Interleukin-6, its -174 G>C Polymorphism and C-Reactive Protein in Idiopathic Cardiac Arrhythmias in Children. Adv Med Sci. 2013;58(2):320-5. doi: 10.2478/ams-2013-0003.
7. Seo IH, Lee YJ. Usefulness of Complete Blood Count (CBC) to Assess Cardiovascular and Metabolic Diseases in Clinical Settings: A Comprehensive Literature Review. Biomedicines. 2022;10(11):2697. doi: 10.3390/biomedicines10112697.
8. Aydın M, Yıldız A, Yüksel M, Polat N, Aktan A, İslamoğlu Y. Assessment of the Neutrophil/Lymphocyte Ratio in Patients with Supraventricular Tachycardia. Anatol J Cardiol. 2016;16(1):29-33. doi: 10.5152/akd.2015.5927.



This is an open-access article distributed under the terms of the Creative Commons Attribution License